

WHAT YOU MAY EXPERIENCE WHEN TERRIBLE THINGS HAPPEN

There are a wide variety of positive and negative reactions that survivors can experience during and immediately after a trauma. These include:

| DOMAIN | NEGATIVE REACTIONS | POSITIVE REACTIONS |
|---------------|-------------------------------------|----------------------------------|
| Cognitive | Confusion, disorientation, worry, | Determination and resolve, |
| | intrusive thoughts and images, | sharper perception, courage, |
| | self-blame | optimism, faith |
| Emotional | Shock, sorry, grief, sadness, fear, | Social connectedness, altruistic |
| | anger, numb, irritability, guilt, | helping behaviors |
| | and shame | |
| Social | Extreme withdrawal, | Social connectedness, altruistic |
| | interpersonal conflict | helping behaviors |
| Psychological | Fatigue, headache, muscle | Alertness, readiness to respond, |
| | tension, stomachache, increased | increased energy |
| | heart rate, exaggerated startle | |
| | response, difficulties sleeping | |

INTRUSIVE REACTIONS

- Distressing thoughts or images of the event while awake or dreaming
- Upsetting emotional or physical reactions to reminders of the experience
- Feeling like the experience is happening all over again (flashback)

AVOIDANCE AND WITHDRAWAL REACTIONS

- Avoid talking, thinking, and having feelings about the traumatic event
- Avoid reminders of the event (places and people connected to what happened)
- Restricted emotions; feeling numb.
- Feelings of detachment and estrangement from others; social withdrawal
- Loss of interest in usually pleasurable activities

PHYSICAL AROUSAL REACTION

- Constantly being "on the lookout" for danger, startling easily, or being jumpy
- Irritability or outburst of anger, feeling "on edge"
- Difficulty falling or staying asleep, problems concentrating or paying attention

REACTIONS TO TRAUMA AND LOSS REMINDERS

- Reactions to places, people, sights, sounds, smells and feelings that are reminders of the disaster
- Reminders can bring on distressing mental images, thoughts, and emotional/physical reactions
- Common examples include sudden loud noises, sirens, locations where the disaster occurred, seeing
 people with disabilities, funerals, anniversaries of the disaster and television/radio news about the
 disaster

POSITIVE CHANGES IN PRIORITIES, WORLDVIEW AND EXPECTATIONS

- Enhanced appreciation that family and friends are precious and important
- Meeting the challenge of addressing difficulties (by taking positive action steps changing the focus of thoughts, using humor, acceptance)

- Shifting expectations about what to expect from day to day and about what is considered a "good day".
 Shifting priorities to focus more on quality time with family or friends
- Increased commitment to self, family friends and spiritual/religious faith

WHEN A LOVED ONE DIES, COMMON REACTIONS INCLUDE:

- Feeling confused, numb, disbelief, bewildered, or lost
- Feeling angry at the person who died or at people considered responsible for the death
- Strong physical reactions such as nausea, fatigue, shakiness, and muscle weakness
- Feeling guilty for still being alive
- Intense emotions such as extreme sadness, anger, or fear
- Increased risk for physical illness and injury
- Decreased productivity or difficulty making decisions
- Having thoughts about the person who died, even when you don't want to
- · Longing, missing, and wanting to search for the person who died
- Children are particularly likely to worry that they or a parent might die
- Children may become anxious when separated from caregivers or other loved ones

WHAT HELPS

- Talking to another person for support or spending time with others
- Engaging in positive distracting activities (sports, hobbies, reading)
- Getting adequate rest and eating healthy meals
- Trying to maintain a normal schedule
- Scheduling pleasant activities
- Taking breaks
- · Reminiscing about a loved one who has died
- Focusing on something practical that you can do right now to manage the situation better
- Using relaxation methods (breathing exercises, meditation, calming, self-talk, soothing music)
- Participating in a support group
- Exercising in moderation
- Keeping a journal
- Seeking counseling

WHAT DOESN'T HELP

- Using alcohol or drugs to cope
- Extreme avoidance of thinking or talking about the event or a death of a loved one
- Violence of conflict
- Overeating or failing to eat
- Excessive TV or computer games
- Blaming others
- Working too much
- Extreme withdrawal from family or friends
- Not taking care of yourself
- Doing risky things (driving recklessly, substance abuse, not taking adequate precautions)
- Withdrawing from pleasant activities

CRISIS RESOURCES

- Crisis Text Line: Text SUPPORT to 741-741
- National Alliance on Mental Illness (NAMI): 800-950-6264
- National Suicide Prevention Lifeline: 1-800-273-TALK (8255)
- Trevor Project (for LGBTQ youth): 1-866-488-7386 or thetrevorproject.org
- Wake Forest Counseling Center 24/7 Call Service: 336-758-5273
- 758-CARE: 336-758-2273