

Dear Allergist,

Wake Forest University Deacon Health looks forward to working with you and your patient. To help us better serve your patient, and to maximize safety, we require the attached PHYSICIAN ORDER FOR ALLERGY

IMMUNOTHERAPY be completed and signed for each allergy serum before we can continue allergy injections. This form will decrease the chance of miscommunication and resultant allergy administration errors.

Please note:

- Patient's initial injection(s) must be performed at an Allergist's office.
- Each vial must be clearly labeled with the patient's name, dilution and expiration date.
- Each Vial must list all materials in the serum.
- No expired serum will be administered. Exceptions may be made if explicit instructions are provided on the order form (i.e. Okay to use serum one month beyond expiration date without need for further consultation with my office)
- New serum vials to be sent directly to the patient, NOT Deacon Health.
- For any systemic reactions, Epinephrine (1:1000) is given IM.
- Patients must also have an EpiPen in their possession at each appointment, in case of a delayed reaction after leaving the office. Please provide your patient with a prescription for the auto-injector, as students who are unable to produce an EpiPen at the time of the appointment will not be able to receive their injection.
- If hives or itching of skin/ mucosal membranes or runny nose are present, Diphenhydramine 50 mg is to be administered IM.
- The allergist office will be notified of the event and a request for further instructions and orders.
- No injections will be given if the patient is ill, febrile, wheezing, has an upper respiratory infection, or has hives or an unknown rash.
- Allergy injections will only be administered in Deacon Health when a provider is in the clinic. Rarely, a physician
 will not be on site, and advanced practice providers will be available to assist your patient with any concerns that
 occur with administration of the serum.

Exclusion criteria

- Patients on a beta-blocker or MAOI
- Patients receiving venom immunotherapy, mixed vespid and whole body extracts will not be administered by Deacon Health Staff.
- Deacon Health reserves the right to discontinue this service and refer back to you for management should any safety concerns develop while under care.

Physician Order for Allergy Immunotherapy

For your patient's safety and to facilitate the transfer of allergy treatment to our clinic, this form must be completed to provide standardization and prevent errors. Failure to complete this form may delay or prevent the patient from utilizing our services. This form can be delivered by the patient, mailed, or faxed (see address and fax below).

Patient Name:	Date of Birth:
Physician:	Practice Name:
Office Phone:	Fax:
Pre-Injection Checklist	
Does your patient have a history of Asth	ma? YES □ NO □
History of Anaphylaxis? YES □ NO □	
Does your patient use antihistamine prior	r to receiving allergy injections? YES \(\sigma\) NO \(\sigma\)
Do you require a Peak Flow prior to inje	ctions? YES \(\sigma \) NO \(\sigma \)
○ If YES , Peak flow must be >	L/min
A mandatory wait time will be enforced	after injection.
o Do you require a 20 □ or 30 □ r	ninute wait?

Allergy Vials

Vial	Vial Contents	Last Dose Given	Dilution of Vial	Date of Last Dose
Ex: Vial A	Cat, Dog, Grass	0.3	1:100	5/1/2022

Injection Schedule

	Frequency of Injections		
Build Up	Every	days.	
Maintenance	Every	days or	weeks.

Dilution					
Vial Cap Color					
Expiration Date(s)	//	//	//	//	//
	ml	ml	ml	ml	ml
	ml	ml	ml	ml	ml
	ml	ml	ml	ml	ml
	ml	ml	ml	ml	ml
	ml	ml	ml	ml	ml
	ml	ml	ml	ml	ml
	Go to Next Dilution	ml	ml	ml	ml
		Go to Next Dilution	ml	ml	ml
			Go to Next Dilution	ml	ml
				Go to Next Dilution	ml
					ml

Management of Missed Injections: (According to # of days from LAST injection)

During Build-Up Phase	After Reaching Maintenance		
to days – continues as scheduled	to weeks – give same maintenance dose		
to days – repeat previous dose	to weeks – reduce previous dose by ml		
to days – reduce previous dose by ml	to weeks – reduce previous dose by ml		
to days – reduce previous dose by ml	Over weeks – contact office for instructions		
Over days – contact office for instructions			

Reactions

At next visit:	Proceed with next dose if swelling is < mm				
	Repeat next dose if swelling is >	mm and <	mm		
	Reduce next dose by	ml if swelling is >	mm		
	Call the office if >	_ mm or systemic reaction			
Other Instruc	etions				
Physician Sign	nature:		Date:		
Office Addres	S:				

Wake Forest University Deacon Health clinical staff will call your office for any clarifications in orders prior to giving injections.